

# Fluid and particle passage in three duiker species

Marcus Clauss · Nicola Lunt · Sylvia Ortmann ·  
Amy Plowman · Daryl Codron · Jürgen Hummel

Received: 7 April 2010 / Revised: 17 May 2010 / Accepted: 23 May 2010 / Published online: 8 June 2010  
© Springer-Verlag 2010

**Abstract** Ruminants are characterised by two different types of reticulorumen (RR) physiology. ‘Cattle-type’ ruminants have, amongst other features such as RR contents stratification and a heterogenous intraruminal papillation, a distinct difference between the mean retention time (MRT) of small particles and fluids (the ratio is called the selectivity factor, SF). ‘Moose-type’ ruminants have RR contents that are less stratified, a more homogenous intraruminal papillation and low SFs, indicating less difference in the MRT of small particles and fluids. To date, physiological data indicating a ‘moose-type’ physiology have only been measured in giraffids and Odocoilean cervids, raising the

question whether it is limited to these taxonomic groups only. Here, we measured MRTs of fluids and particles in five duikers (Bovidae, Cephalophinae) from three species (*Sylvicapra grimmia*, *Cephalophus monticola* and *Cephalophus sylvicultor*) and found SFs in the RR of  $1.27 \pm 0.18$ —well within the range of these other browsers. These results are the first physiological indication that a ‘moose-type’ physiology may also occur in bovid species and thus might represent a true convergent adaptation.

**Keywords** Stratification · Rumen physiology · Particle retention · Browser · Grazer

Communicated by C. Gortázar

M. Clauss (✉) · D. Codron  
Clinic for Zoo Animals, Exotic Pets and Wildlife,  
Vetsuisse Faculty, University of Zurich,  
Winterthurerstr. 260,  
8057 Zurich, Switzerland  
e-mail: mclauss@vetclinics.uzh.ch

N. Lunt  
Antelope Project, Marwell Zimbabwe Trust,  
Box 3863, Bulawayo, Zimbabwe

S. Ortmann  
Leibniz Institute for Zoo and Wildlife Research (IZW),  
Alfred-Kowalke-Str. 17,  
10315 Berlin, Germany

A. Plowman  
Whitley Wildlife Conservation Trust,  
Paignton Zoo, Totnes Road, Paignton,  
Devon TQ4 7EU, UK

J. Hummel  
Institute of Animal Science, University of Bonn,  
Endenicher Allee 15,  
53115 Bonn, Germany

## Introduction

One of the most prominent differences in rumen physiology between different ruminant species is the pattern of fluid vs. particle passage. In many ruminants, fluids pass from the rumen well before small particles, whereas in other species, fluids and particles move almost together. It has been suggested that this difference reflects feeding habits of species, with grazers having a very distinct difference between fluid and particle passage and browsers with a lesser difference (Clauss and Lechner-Doll 2001; Clauss et al. 2006b). More recently, we suggested that ruminants of different physiology should not be defined in terms of their natural diet but in terms of their physiology, so that conceptually, physiology and diet are separated and can thus be compared against each other. Thus, the term ‘cattle-type’ ruminant was suggested for ruminants characterised by a distinct difference in fluid and particle passage and the term ‘moose-type’ ruminant for species characterised by a very slight difference between fluid and particle passage (Clauss et al. 2010c).

The ‘cattle-type’ physiology has been observed in ruminants as taxonomically variable as cervids (*Cervus elaphus*; Renecker and Hudson 1990); domestic and wild cattle, goat and sheep (Lechner-Doll et al. 1990; Gross et al. 1996; Behrend et al. 2004; Flores-Miyamoto et al. 2005; Clauss et al. 2006b; Schwarm et al. 2008); antelope (*Addax nasomaculatus*; Hummel et al. 2008); and muskox (*Ovibos moschatus*; Lechner et al. 2010). A ‘moose-type’ physiology has, so far, only been demonstrated in giraffids (*Giraffa camelopardalis* and *Okapia johnstoni*; Clauss et al. 1998; Hummel et al. 2005) and in the Odocoilean cervids moose (*Alces alces*) and roe deer (*Capreolus capreolus*; Renecker and Hudson 1990; Behrend et al. 2004; Lechner et al. 2010), which raises the question if this physiology is limited to these taxonomic groups only.

Duikers consume browse and fruits in the wild (Gagnon and Chew 2000; Wilson 2005). Contrary to common intuition, the wild fruits that are part of the duikers’ diets have very high fibre content and also contain significant levels of secondary plant compounds (Dierenfeld et al. 2002; Molloy and Hart 2002). Duikers are nonetheless well-adapted to digest high-fibre and tannin-containing diets (Shipley and Felicetti 2002). The anatomy of the forestomach of duikers has been described in detail by Hofmann (1973). When compared to other ruminant species, duikers have relatively large salivary glands (Hofmann et al. 2008), a small rumen with thin ruminal pillars (Clauss et al. 2003) and an even papillation (Faurie and Perrin 1995; Clauss et al. 2009), shallow reticular crests (Clauss et al. 2010b) and a small omasum (Clauss et al. 2006a). As these anatomical features are typical for browsing species and also characterise those species in which a ‘moose-type’ physiology has been observed, we hypothesised that fluids and particles would pass without much difference through the digestive tract of duikers.

Ingesta passage has so far been measured in blue duiker (*Cephalophus monticola*; Luginbuhl et al. 1990; Wenninger and Shipley 2000) and Maxwell’s duiker (*Cephalophus maxwellii*; Conklin-Brittain and Dierenfeld 1996). Although the standard set of passage markers (cobalt–ethylenediaminetetraacetic acid (Co–EDTA) for fluids and chromium–mordanted fibres for particles) used to investigate wild ruminants in the studies listed above was applied in blue duiker (Luginbuhl et al. 1990; Wenninger and Shipley 2000), the pattern of fluid vs. particle retention was not the focus of, and was not discussed in, these previous studies.

## Materials and methods

We used five duikers from three species—grey duiker (*Sylvicapra grimmia*), blue duiker (*C. monticola*) and yellow-backed duiker (*Cephalophus sylvicultor*; Table 1)—

kept at the Dambari Field Station of the Marwell Zimbabwe trust. Animals were kept individually in vegetated enclosures that they had inhabited for more than 1 year. The husbandry and diets of the animals at Dambari Field Station were described by Plowman (2002). In short, duikers received a diet of mixed domestic vegetables (gem squash, butternut, carrot, pumpkin and green banana; 100, 250 and 500 g fresh weight per blue, grey and yellow-backed duiker, respectively), game cubes (National Foods, Harare, Zimbabwe; 150, 450 and 1,000 g as fed) and a variety of browse species indigenous to the area at approximately 1600 hours daily; water was available ad libitum. Uneaten feed was removed by 0800 hours the following morning. Food intake was not recorded throughout the study, as measuring consumption of vegetation growing in enclosures was logistically impractical.

The markers cobalt–EDTA and chromium(Cr)–mordanted fibre (<2 mm) were prepared according to Udén et al. (1980). Animals were fed the markers with a small quantity of vegetables or molasses; the mixture was removed 1 h later and the usual diet was provided. Pens were checked regularly at hourly intervals for the first 48 h and at increasing intervals until approximately 130 h after marker application, and faeces defecated within the respective intervals were sampled and dried at 100°C to constant mass. A faecal sample taken from the animals prior to marker dosage was used as baseline value. Samples were treated and analysed as described by Behrend et al. (2004).

The mean retention time for the whole gastrointestinal tract (MRT GIT) was calculated according to Thielemans et al. (1978) as

$$\text{MRT GIT} = \frac{\sum t_i C_i dt_i}{\sum C_i dt_i}$$

with  $C_i$  = marker concentration in the faecal samples from the interval represented by time  $t_i$  (hours after marker administration) and  $dt_i$  = the interval (hours) of the respective sample

$$dt_i = \frac{(t_{i+1} - t_i) + (t_i - t_{i-1})}{2}$$

Liquid MRTs for the reticulorumen (MRT RR) were calculated by Grovum and Williams (1973b); this calculation is based on the decrease of the faecal liquid marker concentration  $C_i$  with time according to the equation

$$C_i = a e^{-kt_i} \text{ or } \ln C_i = -k t_i + b$$

$\text{MRT}_{\text{fluidRR}}$  then is  $k^{-1}$ . Because markers can be assumed to move in parallel in the distal gastrointestinal tract of ruminants (empirically confirmed by Grovum and Williams 1973a; Kaske and Groth 1997; Mambrini and Peyraud 1997; Wylie et al. 2000),  $\text{MRT}_{\text{particlesRR}}$  was calculated according to Lechner-Doll et al. (1990) by

**Table 1** Animals used in this study, estimated body mass (BM, kg), age (years), mean retention times (MRT, h) in the whole gastrointestinal tract (GIT) and the reticulorumen (RR) for fluids and particles (<2 mm) and selectivity factors (SF; ratio of particle–fluid MRT)

Species	Sex	BM	Age	MRT <sub>GIT</sub>		SF <sub>GIT</sub>	MRT <sub>RR</sub>		SF <sub>RR</sub>
				Fluid	Particles		Fluid	Particles	
<i>S. grimmia</i>	m	13	1	27	27	1.02	21	21	1.02
	f	13	13	23	28	1.24	17	22	1.32
<i>C. monticola</i>	f	4	7	26	35	1.33	20	28	1.44
<i>C. sylvicultor</i>	m	65	12	39	43	1.10	31	35	1.13
	f	65	11	30	38	1.26	19	27	1.41


assuming that  $MRT_{\text{fluid}}^{\text{distal}} = MRT_{\text{fluid}}^{\text{GIT}} - MRT_{\text{fluid}}^{\text{RR}}$  and  $MRT_{\text{particle}}^{\text{distal}} = MRT_{\text{fluid}}^{\text{distal}}$ ; hence,  $MRT_{\text{particles}}^{\text{RR}} = MRT_{\text{particles}}^{\text{GIT}} - MRT_{\text{particles}}^{\text{distal}}$ . Selectivity factors (SFs) according to Lechner-Doll et al. (1990) were calculated for GIT or RR as  $MRT_{\text{particles}}/MRT_{\text{fluid}}$ .

Because digesta passage patterns probably correlate with intraruminal papillation patterns (Clauss et al. 2009), the results of this study were added to a corresponding data collection. The intraruminal papillation pattern is expressed as the proportion of surface enlargement at the site of the smallest surface enlargement factor (SEF)—the dorsal rumen—of the surface enlargement at the site of the highest SEF—the atrium.

Comparisons were made with literature data, and statistical evaluations were performed by regression analysis using PASW 18.0 (SPSS Inc., Chicago, IL, USA).

## Results


Marker excretion curves in the duiker showed very similar patterns for fluids and particles (Fig. 1).  $MRT_{\text{fluid}}^{\text{GIT}}$  and  $MRT_{\text{particle}}^{\text{GIT}}$  ranged from 23 to 39 h and from 27 to 43 h, respectively, with a mean  $SF_{\text{GIT}}$  of  $1.19 \pm 0.13$  h.  $MRT_{\text{fluid}}^{\text{RR}}$  and  $MRT_{\text{particle}}^{\text{RR}}$  ranged from 17 to 31 h and from 21 to 35 h, respectively, with a mean  $SF_{\text{RR}}$  of  $1.27 \pm 0.18$  h. When combining data of body mass (BM) and  $MRT_{\text{particle}}^{\text{GIT}}$  from this study with literature data, there was no significant allometric relationship between the two parameters (Fig. 2).


**Fig. 1** Marker excretion patterns for fluids (Co) and particles (Cr) in a grey duiker (*S. grimmia*)

Compared to results on fluid and particle retention in the RR of other ruminants, duikers fit the pattern of other ‘moose-type’ ruminants (Fig. 3). When combining data on rumen papillation and the  $SF_{\text{RR}}$  for different species (Fig. 4), there was a significant negative linear correlation ( $R = -0.645$ ,  $p = 0.017$ ,  $n = 13$ ;  $SF_{\text{RR}} = 2.39 - 0.013 \text{ SEF}$ ), indicating that larger differences in papillation within the rumen are related to larger differences between fluid and particle passage from the rumen; a better fit to the data was achieved using a logarithmic regression ( $R = 0.781$ ,  $p = 0.002$ ,  $n = 13$ ;  $SF_{\text{RR}} = 3.94 - 0.589 \ln(\text{SEF})$ ).

## Discussion

To our knowledge, the results of this study represent the first physiological indication of a ‘moose-type’ physiology in bovid species. So far, bovid species investigated had always had a distinct difference between fluid and particle passage, i.e. they belonged to the ‘cattle-type’ physiology (anoa *Bubalus depressicornis*, Flores-Miyamoto et al. 2005; domestic cattle, sheep and goats, ibex *Capra ibex* and mouflon *Ovis ammon musimon*, reviewed in Clauss et al. 2006b; addax *Addax nasomaculatus*, Hummel et al. 2008; banteng *Bos javanicus*, Schwarm et al. 2008 and muskoxen *Ovibos moschatus*, Lechner et al. 2010).

**Fig. 2** Body mass and the mean retention time (MRT) of particles in the gastrointestinal tract (GIT) of different duiker species; data from this study (grey circles) and the literature (black circles; Luginbuhl et al. 1990; Conklin-Brittain and Dierenfeld 1996; Wenninger and Shipley 2000)



**Fig. 3** Relationship of the mean retention time (*MRT*) of fluids and small particles in the reticulorumen (*RR*) of various ruminant species classified as ‘cattle-type’ or ‘moose-type’ (from Clauss et al. 2010c) with data for duikers added from this study


The duikers had very low  $SF_{GIT}$  ( $1.19 \pm 0.13$ ) and  $SF_{RR}$  ( $1.27 \pm 0.18$ ) within the typical ‘browser’ range of  $SF_{GIT}$  1.2–1.3 and  $SF_{RR}$  1.1–1.8, respectively (Hummel et al. 2005). Our findings are in accord with a  $SF_{GIT}$  of 1.29 calculated from data for Co–EDTA and Cr-mordanted fibres in blue duikers from Wenninger and Shipley (2000; in that study, however, these two markers were not fed simultaneously); they do not match the  $SF_{GIT}$  of 1.52–1.81 calculated from data for Co–EDTA and Cr-mordanted fibres in blue duikers from Luginbuhl et al. (1990; no particle size given for mordanted fibres, method of calculation of *MRT* not explained). Our study indicates that a strategy of little differentiation between fluid and particle passage from the *RR* is not restricted to giraffids and Odocoilean cervids, can be adopted by bovid species as well and might therefore represent a true convergent adaptation (Fig. 3).

Some limitations of this study need to be mentioned. Apart from the small sample size, the food intake of the duikers could not be recorded for logistic reasons. Food intake is one of the major factors influencing *MRT* (Clauss et al. 2007) and has been shown to also influence *MRT* in duikers; however, in ruminants, food intake does not appear to notably influence *SF*, i.e. the ratio of  $MRT_{particle}/MRT_{fluid}$  (Schwarm et al. 2009), which was the main target of this study. Similarly, diet does not appear to have a major influence on *SF* (cf. Renecker and Hudson 1990); results from this study must therefore be considered meaningful even if not measured on the natural diet of the species.

Wenninger and Shipley (2000) collated literature data that showed that within ruminants of a body mass below 100 kg, there was no indication for an increase in  $MRT_{part,GIT}$  with body mass. Combining literature data and data from this study, there was no significant correlation between *BM* and  $MRT_{part,GIT}$ , either (Fig. 2). Clauss et al. (2007) had already refuted such a relationship based on empirical data, although

theoretical considerations support it (Demment and Van Soest 1985). As in the larger dataset in Clauss et al. (2007), this is mainly due to the fact that even in small species, like the smaller duikers, comparatively long *MRT*s are achieved. Other authors have commented before on the high-fibre digestibilities achieved by duikers (Hart 1986; Conklin-Brittain and Dierenfeld 1996; Wenninger and Shipley 2000; Pérez-Barberia et al. 2004), most likely due to the long *MRT*s in these species. The capacity for long *MRT*s may well be an adaptation to the high level of fibre in the natural diet of duikers as described in the “Introduction”.

Anatomical characteristics are often used to differentiate functional guilds of ruminants (Hofmann 1989; Mendoza and Palmqvist 2006). With respect to *RR* physiology and *RR* contents stratification, the intraruminal papillation pattern is an important proxy (Clauss et al. 2009), distinguishing ‘moose-type’ from ‘cattle-type’ species (Codron and Clauss, subm.). However, such interpretations will often have to be tempered by the fact that the relationship between form (an anatomical proxy, such as the papillation pattern) and function (a physiological proxy, such as the *SF*) is nonlinear. In the comparison of the intraruminal papillation pattern with *SF* measurements in Fig. 4, there is no clear correlation above or below the thresholds of an  $SF_{RR}$  of 2.0 and a dorsal surface enlargement factor of 30% of the *SEF* in the *Atrium ruminis*. A similar trend was observed in the correlation of



**Fig. 4** Relationship between the selectivity factor (*SF*; ratio of the mean retention time of particles in the reticulorumen (*RR*) to that of fluids—a physiological indicator of *RR* contents stratification) and the intraruminal papillation pattern (an anatomical indicator of *RR* contents stratification). The intraruminal papillation pattern is expressed as the proportion of surface enlargement at the site of the smallest surface enlargement factor [*SEF*—the dorsal rumen—of the surface enlargement at the site of the highest *SEF*—the atrium. Original data collection from Clauss et al. (2009), with additional results from recent studies on muskoxen (*O. moschatus*) and reindeer (*Rangifer tarandus*; Clauss et al. 2010a; Lechner et al. 2010) and with data on duiker from this study (*grey circles*; *MRT* data from this study for *S. grimmia* and *C. monticola*, linked to papillation data from Clauss et al. 2009 on *S. grimmia* and *Cephalophus harveyi*). Logarithmic regression is described in “Results”



intraruminal papillation with another measure of RR contents stratification—the difference in dry matter concentration between the dorsal and the ventral RR content (Codron and Clauss, *subm.*). Similar nonlinear or threshold-dependent effects could be expected for different morphological measurements.

To conclude, the results of our trials indicate that similar to other ruminants depending on diets with consistent amounts of tannins, duikers are characterised by a low difference in the MRT of small particles and fluids, which indicates a low degree of RR content stratification and thus fits the observed intraruminal papillation patterns. Such a physiology is probably linked to other features common to these and similar animals, for example large salivary glands that potentially produce tannin-binding proteins and hence a more viscous saliva (Hofmann et al. 2008). While proximate causes for the observed variation in particle and fluid retention remain to be investigated, comparative retention measurements are an important addition to our understanding of ruminant physiology.

**Acknowledgements** We thank the Director and staff at Marwell Zimbabwe Trust for assistance with sample collection and Heidrun Barleben from the IZW for her engaged support in the lab. Whitley Wildlife Conservation Trust (UK) and Marwell Wildlife (UK) funded duiker husbandry.

## References

- Behrend A, Lechner-Doll M, Streich WJ, Clauss M (2004) Seasonal faecal excretion, gut fill, liquid and particle marker retention in mouflon (*Ovis ammon musimon*), and a comparison with roe deer (*Capreolus capreolus*). *Acta Theriol* 49:503–515
- Clauss M, Lechner-Doll M (2001) Differences in selective reticulo-ruminal particle retention as a key factor in ruminant diversification. *Oecologia* 129:321–327
- Clauss M, Deutsch A, Lechner-Doll M, Flach EJ, Tack C (1998) Passage rate of fluid and particle phase in captive giraffe (*Giraffa camelopardalis*). *Adv Ethol* 33:98
- Clauss M, Lechner-Doll M, Streich WJ (2003) Ruminant diversification as an adaptation to the physicochemical characteristics of forage. A reevaluation of an old debate and a new hypothesis. *Oikos* 102:253–262
- Clauss M, Hofmann RR, Hummel J, Adamczewski J, Nygren K, Pitra C, Streich WJ, Reese S (2006a) The macroscopic anatomy of the omasum of free-ranging moose (*Alces alces*) and muskoxen (*Ovibos moschatus*) and a comparison of the omasal laminal surface area in 34 ruminant species. *J Zool Lond* 270:346–358
- Clauss M, Hummel J, Streich WJ (2006b) The dissociation of the fluid and particle phase in the forestomach as a physiological characteristic of large grazing ruminants: an evaluation of available, comparable ruminant passage data. *Eur J Wildl Res* 52:88–98
- Clauss M, Schwarm A, Ortmann S, Streich WJ, Hummel J (2007) A case of non-scaling in mammalian physiology? Body size, digestive capacity, food intake, and ingesta passage in mammalian herbivores. *Comp Biochem Physiol A* 148:249–265
- Clauss M, Hofmann RR, Fickel J, Streich WJ, Hummel J (2009) The intraruminal papillation gradient in wild ruminants of different feeding types: implications for rumen physiology. *J Morphol* 270:929–942
- Clauss M, Adamczewski J, Hofmann RR (2010a) Surface enlargement in the rumen of free-ranging muskoxen (*Ovibos moschatus*). *Eur J Wildl Res* 56:181–185
- Clauss M, Hofmann RR, Streich WJ, Fickel J, Hummel J (2010b) Convergence in the macroscopic anatomy of the reticulum in wild ruminant species of different feeding types and a new resulting hypothesis on reticular function. *J Zool Lond* 281:26–38
- Clauss M, Hume ID, Hummel J (2010c) Evolutionary adaptations of ruminants and their potential relevance for modern production systems. *Animal* 4:979–992
- Codron D, Clauss M (2010) Differences in rumen fluid concentrations support stratification as a constraint to browsing and grazing in ruminants (submitted)
- Conklin-Brittain NL, Dierenfeld ES (1996) Small ruminants: digestive capacity differences among four species weighing less than 20 kg. *Zoo Biol* 15:481–490
- Demment MW, Van Soest PJ (1985) A nutritional explanation for body size patterns of ruminant and nonruminant herbivores. *Am Nat* 125:641–672
- Dierenfeld ES, Mueller PJ, Hall MB (2002) Duikers: native food composition, micronutrient assessment, and implications for improving captive diets. *Zoo Biol* 21:185–196
- Faurie AS, Perrin MR (1995) Rumen morphology and volatile fatty acid production in the blue duiker and the red duiker. *Mamm Biol* 60:73–84
- Flores-Miyamoto K, Clauss M, Ortmann S, Sainsbury AW (2005) The nutrition of captive lowland anoa (*Bubalus depressicornis*): a study on ingesta passage, intake, digestibility, and a diet survey. *Zoo Biol* 24:125–134
- Gagnon M, Chew AE (2000) Dietary preferences in extant African Bovidae. *J Mammal* 81:490–511
- Gross JE, Alkon PU, Demment MW (1996) Nutritional ecology of dimorphic herbivores: digestion and passage rates in *Nubian ibex*. *Oecologia* 107:170–178
- Grovum WL, Williams VJ (1973a) Rate of passage of digesta in sheep: 4. Passage of marker through the alimentary tract and the biological relevance of rate-constants derived from the changes in concentration of marker in faeces. *Br J Nutr* 30:313–329
- Grovum WL, Williams VJ (1973b) Rate of passage of digesta in sheep. 3. Differential rates of passage of water and dry matter from the reticulo-rumen, abomasum and caecum and proximal colon. *Br J Nutr* 30:231–240
- Hart JA (1986) Comparative dietary ecology of a community of frugivorous forest ungulates in Zaire. Ph.D. Dissertation, Michigan State University, East Lansing, MI
- Hofmann RR (1973) The ruminant stomach. East African Literature Bureau, Nairobi
- Hofmann RR (1989) Evolutionary steps of ecophysiological adaptation and diversification of ruminants: a comparative view of their digestive system. *Oecologia* 78:443–457
- Hofmann RR, Streich WJ, Fickel J, Hummel J, Clauss M (2008) Convergent evolution in feeding types: salivary gland mass differences in wild ruminant species. *J Morphol* 269:240–257
- Hummel J, Clauss M, Zimmermann W, Johanson K, Norgaard C, Pfeffer E (2005) Fluid and particle retention in captive okapi (*Okapia johnstoni*). *Comp Biochem Physiol A* 140:436–444
- Hummel J, Steuer P, Südekum K-H, Hammer S, Hammer C, Streich WJ, Clauss M (2008) Fluid and particle retention in the digestive tract of the addax antelope (*Addax nasomaculatus*)—adaptations of a grazing desert ruminant. *Comp Biochem Physiol A* 149:142–149
- Kaske M, Groth A (1997) Changes in factors affecting the rate of digesta passage through pregnancy and lactation in sheep fed on hay. *Reprod Nutr Dev* 37:573–588

- Lechner I, Barboza P, Collins W, Fritz J, Günther D, Hattendorf B, Hummel J, Südekum K-H, Clauss M (2010) Differential passage of fluids and different-sized particles in fistulated oxen (*Bos primigenius* f. *taurus*), muskoxen (*Ovibos moschatus*), reindeer (*Rangifer tarandus*) and moose (*Alces alces*): rumen particle size discrimination is independent from contents stratification. *Comp Biochem Physiol A* 155:211–222
- Lechner-Doll M, Rutagwenda T, Schwartz HJ, Schultka W, von Engelhardt W (1990) Seasonal changes of ingesta mean retention time and forestomach fluid volume in indigenous camels, cattle, sheep and goats grazing in a thornbush savanna pasture in Kenya. *J Agric Sci (Cambridge)* 115:409–420
- Luginbuhl JM, Pond KR, Burns JC (1990) Physical limits to ruminal escape: experiences with the blue duiker. In: Oftedal OT, Barboza PS (eds) Digestive strategies of animals, a symposium. National Zoological Park, Smithsonian Institution, Washington DC, pp 10–11
- Mambrini M, Peyraud JL (1997) Retention time of feed particles and liquids in the stomachs and intestines of dairy cows. Direct measurement and calculation based on fecal collection. *Reprod Nutr Dev* 37:427–442
- Mendoza M, Palmqvist P (2006) Characterizing adaptive morphological patterns related to diet in Bovidae. *Acta Zoologica Sinica* 52:988–1008
- Molloy L, Hart JA (2002) Duiker food selection: palatability trials using natural foods in the Ituri Forest, Democratic Republic of Congo. *Zoo Biol* 21:149–159
- Pérez-Barbería FJ, Elston DA, Gordon IJ, Illius AW (2004) The evolution of phylogenetic differences in the efficiency of digestion in ruminants. *Proc -Royal Soc B* 271:1081–1090
- Plowman AB (2002) Nutrient intake and apparent digestibility of diets consumed by captive duikers at the Dambari Field Station, Zimbabwe. *Zoo Biol* 21:135–147
- Renecker LA, Hudson RJ (1990) Digestive kinetics of moose, wapiti and cattle. *Anim Prod* 50:51–61
- Schwarm A, Ortmann S, Wolf C, Streich WJ, Clauss M (2008) Excretion patterns of fluids and particle passage markers of different size in banteng (*Bos javanicus*) and pygmy hippopotamus (*Hexaprotodon liberiensis*): two functionally different foregut fermenters. *Comp Biochem Physiol A* 150:32–39
- Schwarm A, Ortmann S, Wolf C, Streich WJ, Clauss M (2009) Passage marker excretion in red kangaroo (*Macropus rufus*), collared peccary (*Pecari tajacu*) and colobine monkeys (*Colobus angolensis*, *C. polykomos*, *Trachypithecus johnii*). *J Exp Zool* 311:647–661
- Shipley LA, Felicetti L (2002) Fiber digestibility and nitrogen requirements of blue duikers. *Zoo Biol* 21:123–134
- Thielemans MF, Francois E, Bodart C, Thewis A (1978) Mesure du transit gastrointestinal chez le porc à l'aide des radiolanthanides. Comparaison avec le mouton. *Ann Biol Anim Biochim Biophys* 18:237–247
- Udén P, Colucci PE, Van Soest PJ (1980) Investigation of chromium, cerium and cobalt as markers in digesta rates of passage studies. *J Sci Food Agric* 31:625–632
- Wenninger PS, Shipley LA (2000) Harvesting, rumination, digestion, and passage of fruit and leaf diets by a small ruminant, the blue duiker. *Oecologia* 123:466–474
- Wilson VJ (2005) Duikers of Africa. Chipangali Wildlife Trust, Duiker Research and Breeding Centre, Zimbabwe
- Wylie MJ, Ellis WC, Matis JH, Bailey EM, James WD, Beever DE (2000) The flow of forage particles and solutes through segments of the digestive tracts of cattle. *Br J Nutr* 83:295–306